

Immunoglobulin with High-Titer *In Vitro* Cross-Neutralizing Hepatitis C Virus Antibodies Passively Protects Chimpanzees from Homologous, but Not Heterologous, Challenge

Jens Bukh,^{a,b} Ronald E. Engle,^b Kristina Faulk,^b Richard Y. Wang,^c Patrizia Farci,^b Harvey J. Alter,^c Robert H. Purcell^b

Copenhagen Hepatitis C Program (CO-HEP), Department of Infectious Diseases and Clinical Research Centre, Hvidovre Hospital, and Department of Immunology and Microbiology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark^a; Laboratory of Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA^b; Department of Transfusion Medicine, Grant Magnuson Clinical Center, National Institutes of Health, Bethesda, Maryland, USA^c

The importance of neutralizing antibodies (NAbs) in protection against hepatitis C virus (HCV) remains controversial. We infused a chimpanzee with H06 immunoglobulin from a genotype 1a HCV-infected patient and challenged with genotype strains efficiently neutralized by H06 *in vitro*. Genotype 1a NAbs afforded no protection against genotype 4a or 5a. Protection against homologous 1a lasted 18 weeks, but infection emerged when NAb titers waned. However, 6a infection was prevented. The differential *in vivo* neutralization patterns have implications for HCV vaccine development.

Chimpanzees have been essential for hepatitis C virus (HCV) research (1–3). Understanding the role of neutralizing antibodies (NAbs) in preventing HCV infection is important for vaccine efforts against this important pathogen, which annually infects ~4 million people and causes chronic liver disease (4–12). We showed that chronic-phase patient serum/plasma prevented HCV infection of chimpanzees if the virus and anti-HCV antibody were incubated prior to inoculation (13, 14). Others showed that immunoglobulin (IgG) purified from anti-HIV antibody-positive blood donors prevented HCV infection when mixed with virus and then administered to a chimpanzee (15). However, the protective effect of preexisting polyclonal HCV NAbs remains to be determined.

Studies with culture systems using pseudotyped particles (HCVpp) or infectious viruses (chimeric cell culture-derived HCV [HCVcc]) confirmed the existence of NAbs in acute- and chronic-phase patients, with high-level cross-neutralization of heterotypic HCV strains (16–20). However, differences exist in neutralization capacities of chronic-phase samples against HCV variants of the same genotype (21), and there is a constant evolution of HCV variants escaping NAbs (14, 22, 23). Nonetheless, these *in vitro* data suggest that chronic-phase HCV samples might be useful for broad protection against HCV. Effective NAbs could help define critical epitopes for vaccine development (24–27).

We prepared H06 IgG from plasma obtained almost 30 years after disease onset in patient H with persistent HCV infection (28). By infusing H06 IgG with high *in vitro* neutralizing titers into a chimpanzee (CHA5A009) and then challenging with different HCV strains sensitive to neutralization *in vitro* (18, 19), our aim was to test the principle of passive immunoprophylaxis against homologous and heterologous strains *in vivo*.

Animal experimentation and sample collection from chimpanzee CHA5A009 were performed from 2007 to 2008. The housing and care of the chimpanzee met or exceeded all requirements of the National Research Council's 1996 *Guide for the Care and Use of Laboratory Animals*, 7th ed. (National Academies Press, Washington, DC), which were in effect until 2010. The project license (ASP LID 64) and specific protocol (06-C-482) were separately approved by the Institutional Animal Care and Use Com-

mittees of the National Institute of Allergy and Infectious Diseases, NIH, and Bioqual, Inc., the facility housing the chimpanzee. Both facilities were fully accredited by the American Association for Accreditation of Laboratory Animal Care International. Specifically, housing exceeded requirements for square footage and cage height for the chimpanzee's size. Feeding consisted of approved chimpanzee biscuits with supplemental vegetables and fruit and *ad libitum* water. A comprehensive enrichment program was in place and consisted of foods, puzzles, and environmental enrichment, coupled with visual and auditory contact with other chimpanzees. Sample collection was performed under ketamine hydrochloride anesthesia. Euthanasia was not required.

CHA5A009 was infused with H06 IgG intravenously prechallenge (250 mg/kg body weight) and at weeks 2.5 and 3.5 postchallenge (125 mg/kg). Twenty-four hours after the first IgG infusion, CHA5A009 was challenged intravenously with 100 chimpanzee infectious doses of each of the HCV strains (with the genotype in parentheses) H77(1a), ED43(4a), SA13(5a), and HK6a(6a) (29–31), chosen because H77 is the homologous acute-phase patient H strain (31) and chronic-phase patient H sera cross-neutralized all strains in HCVpp and HCVcc assays (18, 19). Based on 5'- untranslated region (UTR)-based TaqMan assay (32), CHA5A009 had serum HCV titers of 2.7 and 4.8 log₁₀ IU/ml at weeks 1 and 2, respectively, and peak titers of 5 to 6 log₁₀ IU/ml at weeks 3 to 11; the animal remained persistently infected (Fig. 1A). 5'-UTR (33) and core-envelope 1 (core-E1) (34) analysis of week 2 viruses,

Received 9 May 2015 Accepted 15 June 2015

Accepted manuscript posted online 17 June 2015

Citation Bukh J, Engle RE, Faulk K, Wang RY, Farci P, Alter HJ, Purcell RH. 2015. Immunoglobulin with high-titer *in vitro* cross-neutralizing hepatitis C virus antibodies passively protects chimpanzees from homologous, but not heterologous, challenge. *J Virol* 89:9128–9132. doi:10.1128/JVI.01194-15.

Editor: M. S. Diamond

Address correspondence to Jens Bukh, jbukh@sund.ku.dk.

Copyright © 2015, American Society for Microbiology. All Rights Reserved. doi:10.1128/JVI.01194-15

TABLE 1 Primers used in the HCV strain-specific RT-nested PCR in this study

Strain ^a and primer type	Primer	Sequence
H77(1a)		
RT	2832R-H77	AAGCGCCCCTAACTTGATGATG
PCR I	2427S-H77	ACTGGACACGGAGGTGGCCGCGTC
	2832R-H77	AAGCGCCCCTAACTTGATGATG
PCR II	2462S-H77	TTGTTCTTGTCGGGTTAATGGCGC
	2645R-H77	GGGTGTACTACACACATGAGTAAG
ED43(4a)		
RT	HCV4aCoseR3591	CCGTGGTAGACGGTCCACATCAC
PCR I	2676F-ED43	AGGGCCGGTCCCAGCTGCT
	HCV4aCoseR3591	CCGTGGTAGACGGTCCACATCAC
PCR II	2839F-ED43	GCACTACAAGTTATGGCTGGCTA
	HCV4aCoseR3318	CACGCAGCGGTGTCAGCGCCCC
SA13(5a)		
RT	917R-SA13	CAGTTGCAGTCCTGCACAACATTA
PCR I	741S-SA13	AGCCCCGAGCCTCGGAGCGGT
	917R-SA13	CAGTTGCAGTCCTGCACAACATTA
PCR II	748S-SA13	AGCCTCGGAGCGGTACCGGCT
	898R-SA13	CATTATGCCGCGAGGGCTAT
HK6a(6a)		
RT	1765R-HK6a	TGGGGCACAAAAGCTCATTGG
PCR I	1405S-HK6a	TGGGGCCAAATAACCTACAAA
	1765R-HK6a	TGGGGCACAAAAGCTCATTGG
PCR II	1408S-HK6a	GGCCAAATAACCTACAAAAGTC
	1762R-HK6a	GGCACAAAAGCTCATTGGCAG

^a Genotypes are given in parentheses.

taken prior to IgG infusion had reciprocal neutralization titers of <50 against all HCVpp strains. Following IgG infusion, CHA5A009 had significant NAb titers against all strains, remaining at ≥ 100 through week 11; at weeks 20 and 24, the titers were ≤ 50 for all strains (Fig. 1B).

Sera from weeks 2 to 56 were tested for H77, ED43, SA13, and HK6a HCV RNA by RT-nested PCR (Fig. 1A), using strain-specific primers (Table 1) (36). The sensitivities of the H77, SA13, and HK6a assays were equivalent to that using 5'-UTR primers when testing dilutions of H77, SA13, and HK6a pools, respectively (30); the 4a assay was suboptimal. Amplicons were in most cases sequenced, confirming strain authenticity. There was no evidence of H77(1a) infection until week 18, when H77 RNA was detected at 2 log₁₀ genome equivalents (GE)/ml (Fig. 1A and C) (36). The H77(1a) emergence followed the decrease in H77 NAb titers to <50 (Fig. 1B). The week 18 H77(1a) envelope proteins, derived from overlapping amplicons (37), had no changes compared to the polyclonal challenge virus (38, 39), indicating that viral emergence was not due to escape mutations. The animal remained H77 infected, with titers of 4 to 5 log₁₀ GE/ml from weeks 20 to 24; titers fluctuated at low levels thereafter (Fig. 1C).

Despite relatively high NAb titers against SA13(5a) (Fig. 1B), there was no apparent *in vivo* protection against this strain, which appeared at 5 to 6 log₁₀ GE/ml in samples from weeks 2 to 11 (Fig. 1C); infection resolved after week 16 (Fig. 1A and C). The envelope sequence of week 2 SA13(5a), derived from overlapping amplicons (29, 37), had no changes compared to the polyclonal challenge virus (29, 40), indicating failure of neutralization.

CHA5A009 became ED43(4a) infected, with sequence-confirmed ED43 RNA positivity from weeks 2 to 18 (Fig. 1A); since we did not develop a 4a-specific assay with optimized sensitivity, titer or outcome analysis was not performed. Finally, the chimpanzee apparently was not infected with HK6a(6a), being HK6a RNA negative from weeks 2 to 56 (Fig. 1A).

This is the first study to address whether polyclonal NABs, given prechallenge, can prevent HCV infection in chimpanzees. While the homologous virus was suppressed initially, there was failure of neutralization against 2 of 3 heterologous HCV strains. Furthermore, the homologous virus emerged following the disappearance of passively administered NABs and led to persistent infection. This supports the observation that polyclonal antibodies given postchallenge could control HCV infection only temporarily in the chimpanzee model (41). The infused NABs might have prevented infection with heterologous 6a virus, and this was the strain previously found to be most efficiently neutralized in HCVcc assays (19, 42).

We previously demonstrated that chronic-phase patient H IgG, infused prechallenge, could control H77 infection in human liver chimeric mice (28, 43). However, the animals could only be followed short term. The reciprocal *in vitro* neutralization titers detected prechallenge and at weeks 4 and 8 in CHA5A009 (Fig. 1B) were similar to those detected prechallenge in human liver chimeric mice (43). Thus, for the homologous strain, it appeared that polyclonal immunoglobulin with *in vitro* neutralizing activity can control viremia *in vivo*. The chronic-phase sera from patient H had high *in vitro* neutralization titers against the heterologous 4a, 5a, and 6a strains (18, 19), and we previously found partial protection in human liver chimeric mice loaded with H06 IgG against ED43(4a) and HK6a(6a) (28). Here we observed reciprocal neutralization titers of >400 prechallenge and at week 4 postchallenge in the H06-infused chimpanzee using the same HCVpp assays for these 3 strains. Yet, the chimpanzee became infected with the 4a and 5a strains and developed acute hepatitis. For the 5a strain, we showed that this could represent failure of neutralization, since the challenge virus without envelope mutations appeared at high titers at week 2. In evaluating the outcome for the 1a, 4a, 5a, and 6a challenges, it should be recognized that the HCVpp neutralization titers in the chimpanzee were lower than those previously detected for the different genotype stains in chronic-phase sera from patient H (18). Furthermore, it must be recognized that our study in a single chimpanzee does not necessarily permit broad conclusions.

In a recent study, prechallenge infusion with 250 mg/kg of a humanized monoclonal anti-E2 prevented infection in an H77-challenged chimpanzee (44). Using polyclonal immunoglobulin with an HCV-specific antibody concentration of <250 mg/kg, we observed control of H77 viremia but not sterilizing immunity, even though we also infused the animal with IgG postchallenge. In our study, viral interference could have influenced the outcome given the simultaneous infections with 4a and 5a strains.

Recent data indicate that the HCV clearance is associated with NABs (23, 45, 46). During chronic infection, HCV persists despite the presence of high-titer NABs, perhaps because it is shielded from neutralization (47–50) or neutralization-resistant mutants are continuously developing (22, 23, 51). It was demonstrated *in vitro* that chronic-phase patient H sera neutralized only variants from time points early in infection, not later variants (22). Our data confirm *in vivo* that chronic-phase patient H immunoglob-

ulin can control the acute-phase virus. We previously showed that different recombinant genotype viruses, lacking HVR1, were efficiently neutralized by H06 IgG, indicating that it targets conserved neutralization epitopes within E1 and/or E2 (42).

This study sheds new light on the effectiveness of NAb in preventing HCV. It is encouraging that the homologous virus apparently could be suppressed long term. However, the lack of protection against heterologous strains highlights significant issues regarding the possibility of developing broadly protective immunoglobulin preparations using patient-derived antibodies alone. More likely, efficient neutralization will require monoclonal antibodies against conserved conformational epitopes and/or combinations of antibodies targeting different HCV epitopes or viral variants (44, 50, 52–54).

ACKNOWLEDGMENTS

We thank Alicia Brockington for technical assistance and Charlene Shaver for animal care.

This study was supported by the Intramural Research Program of the NIAID, NIH, and by NIAID contract NO1-AO-62713, as well as by research grants to J.B. from the Danish Council for Independent Research-Medical Sciences, the Lundbeck Foundation, and the Novo Nordisk Foundation. J.B. is the recipient of an Advanced-Top Researcher grant from the Danish Council for Independent Research (2014) and the 2015 Novo Nordisk Prize.

The authors declare that they have no conflicts of interest to report.

REFERENCES

- Bukh J, Forns X, Emerson SU, Purcell RH. 2001. Studies of hepatitis C virus in chimpanzees and their importance for vaccine development. *Intervirology* 44:132–142. <http://dx.doi.org/10.1159/000050040>.
- Lanford RE, Bigger C, Bassett S, Klimpel G. 2001. The chimpanzee model of hepatitis C virus infections. *ILAR J* 42:117–126. <http://dx.doi.org/10.1093/ilar.42.2.117>.
- Bukh J. 2012. Animal models for the study of hepatitis C virus infection and related liver disease. *Gastroenterology* 142:1279–1287. <http://dx.doi.org/10.1053/j.gastro.2012.02.016>.
- Forns X, Bukh J, Purcell RH. 2002. The challenge of developing a vaccine against hepatitis C virus. *J Hepatol* 37:684–695. [http://dx.doi.org/10.1016/S0168-8278\(02\)00308-2](http://dx.doi.org/10.1016/S0168-8278(02)00308-2).
- Ball JK, Tarr AW, McKeating JA. 2014. The past, present and future of neutralizing antibodies for hepatitis C virus. *Antiviral Res* 105:100–111. <http://dx.doi.org/10.1016/j.antiviral.2014.02.013>.
- Houghton M, Abrignani S. 2005. Prospects for a vaccine against the hepatitis C virus. *Nature* 436:961–966. <http://dx.doi.org/10.1038/nature04081>.
- Houghton M. 2011. Prospects for prophylactic and therapeutic vaccines against the hepatitis C viruses. *Immunol Rev* 239:99–108. <http://dx.doi.org/10.1111/j.1600-065X.2010.00977.x>.
- Ray R. 2011. Progress toward development of a hepatitis C vaccine with broad shoulders. *Sci Transl Med* 3:94ps33. <http://dx.doi.org/10.1126/scitranslmed.3002772>.
- Honegger JR, Zhou Y, Walker CM. 2014. Will there be a vaccine to prevent HCV infection? *Semin Liver Dis* 34:79–88. <http://dx.doi.org/10.1055/s-0034-1371081>.
- Dahari H, Feinstone SM, Major ME. 2010. Meta-analysis of hepatitis C virus vaccine efficacy in chimpanzees indicates an importance for structural proteins. *Gastroenterology* 139:965–974. <http://dx.doi.org/10.1053/j.gastro.2010.05.077>.
- Alter HJ, Seeff LB. 2000. Recovery, persistence, and sequelae in hepatitis C virus infection: a perspective on long-term outcome. *Semin Liver Dis* 20:17–35. <http://dx.doi.org/10.1055/s-2000-9505>.
- Bukh J, Miller RH, Kew MC, Purcell RH. 1993. Hepatitis C virus RNA in southern African blacks with hepatocellular carcinoma. *Proc Natl Acad Sci U S A* 90:1848–1851. <http://dx.doi.org/10.1073/pnas.90.5.1848>.
- Farci P, Alter HJ, Wong DC, Miller RH, Govindarajan S, Engle R, Shapiro M, Purcell RH. 1994. Prevention of hepatitis C virus infection in chimpanzees after antibody-mediated in vitro neutralization. *Proc Natl Acad Sci U S A* 91:7792–7796. <http://dx.doi.org/10.1073/pnas.91.16.7792>.
- Farci P, Bukh J, Purcell RH. 1997. The quasispecies of hepatitis C virus and the host immune response. *Springer Semin Immunopathol* 19:5–26. <http://dx.doi.org/10.1007/BF00945022>.
- Yu MY, Bartosch B, Zhang P, Guo ZP, Renzi PM, Shen LM, Granier C, Feinstone SM, Cosset FL, Purcell RH. 2004. Neutralizing antibodies to hepatitis C virus (HCV) in immune globulins derived from anti-HCV-positive plasma. *Proc Natl Acad Sci U S A* 101:7705–7710. <http://dx.doi.org/10.1073/pnas.0402458101>.
- Bartosch B, Bukh J, Meunier JC, Granier C, Engle RE, Blackwelder WC, Emerson SU, Cosset FL, Purcell RH. 2003. In vitro assay for neutralizing antibody to hepatitis C virus: evidence for broadly conserved neutralization epitopes. *Proc Natl Acad Sci U S A* 100:14199–14204. <http://dx.doi.org/10.1073/pnas.2335981100>.
- Logvinoff C, Major ME, Oldach D, Heyward S, Talal A, Balfe P, Feinstone SM, Alter H, Rice CM, McKeating JA. 2004. Neutralizing antibody response during acute and chronic hepatitis C virus infection. *Proc Natl Acad Sci U S A* 101:10149–10154. <http://dx.doi.org/10.1073/pnas.0403519101>.
- Meunier JC, Engle RE, Faulk K, Zhao M, Bartosch B, Alter H, Emerson SU, Cosset FL, Purcell RH, Bukh J. 2005. Evidence for cross-genotype neutralization of hepatitis C virus pseudo-particles and enhancement of infectivity by apolipoprotein C1. *Proc Natl Acad Sci U S A* 102:4560–4565. <http://dx.doi.org/10.1073/pnas.0501275102>.
- Scheel TK, Gottwein JM, Jensen TB, Prentoe JC, Hoegh AM, Alter HJ, Eugen-Olsen J, Bukh J. 2008. Development of JFH1-based cell culture systems for hepatitis C virus genotype 4a and evidence for cross-genotype neutralization. *Proc Natl Acad Sci U S A* 105:997–1002. <http://dx.doi.org/10.1073/pnas.0711044105>.
- Gottwein JM, Scheel TK, Jensen TB, Lademann JB, Prentoe JC, Knudsen ML, Hoegh AM, Bukh J. 2009. Development and characterization of hepatitis C virus genotype 1–7 cell culture systems: role of CD81 and scavenger receptor class B type I and effect of antiviral drugs. *Hepatology* 49:364–377. <http://dx.doi.org/10.1002/hep.22673>.
- Tarr AW, Urbanowicz RA, Hamed MR, Albecka A, McClure CP, Brown RJ, Irving WL, Dubuisson J, Ball JK. 2011. Hepatitis C patient-derived glycoproteins exhibit marked differences in susceptibility to serum neutralizing antibodies: genetic subtype defines antigenic but not neutralization serotype. *J Virol* 85:4246–4257. <http://dx.doi.org/10.1128/JVI.01332-10>.
- von Hahn T, Yoon JC, Alter H, Rice CM, Rehermann B, Balfe P, McKeating JA. 2007. Hepatitis C virus continuously escapes from neutralizing antibody and T-cell responses during chronic infection in vivo. *Gastroenterology* 132:667–678. <http://dx.doi.org/10.1053/j.gastro.2006.12.008>.
- Dowd KA, Netski DM, Wang XH, Cox AL, Ray SC. 2009. Selection pressure from neutralizing antibodies drives sequence evolution during acute infection with hepatitis C virus. *Gastroenterology* 136:2377–2386. <http://dx.doi.org/10.1053/j.gastro.2009.02.080>.
- Law JL, Chen C, Wong J, Hockman D, Santer DM, Frey SE, Belshe RB, Wakita T, Bukh J, Jones CT, Rice CM, Abrignani S, Tyrrell DL, Houghton M. 2013. A hepatitis C virus (HCV) vaccine comprising envelope glycoproteins gpE1/gpE2 derived from a single isolate elicits broad cross-genotype neutralizing antibodies in humans. *PLoS One* 8:e59776. <http://dx.doi.org/10.1371/journal.pone.0059776>.
- Meunier JC, Gottwein JM, Houghton M, Russell RS, Emerson SU, Bukh J, Purcell RH. 2011. Vaccine-induced cross-genotype reactive neutralizing antibodies against hepatitis C virus. *J Infect Dis* 204:1186–1190. <http://dx.doi.org/10.1093/infdis/jir511>.
- Ray R, Meyer K, Banerjee A, Basu A, Coates S, Abrignani S, Houghton M, Frey SE, Belshe RB. 2010. Characterization of antibodies induced by vaccination with hepatitis C virus envelope glycoproteins. *J Infect Dis* 202:862–866. <http://dx.doi.org/10.1086/655902>.
- Wong JA, Bhat R, Hockman D, Logan M, Chen C, Levin A, Frey SE, Belshe RB, Tyrrell DL, Law JL, Houghton M. 2014. Recombinant hepatitis C virus envelope glycoprotein vaccine elicits antibodies targeting multiple epitopes on the envelope glycoproteins associated with broad cross-neutralization. *J Virol* 88:14278–14288. <http://dx.doi.org/10.1128/JVI.01911-14>.
- Meuleman P, Bukh J, Verhoye L, Farhoudi A, Vanwolleghem T, Wang RY, Desombere I, Alter H, Purcell RH, Leroux-Roels G. 2011. In vivo evaluation of the cross-genotype neutralizing activity of polyclonal antibodies against hepatitis C virus. *Hepatology* 53:755–762. <http://dx.doi.org/10.1002/hep.24171>.

29. Bukh J, Apgar CL, Engle R, Govindarajan S, Hegerich PA, Tellier R, Wong DC, Elkens R, Kew MC. 1998. Experimental infection of chimpanzees with hepatitis C virus of genotype 5a: genetic analysis of the virus and generation of a standardized challenge pool. *J Infect Dis* 178:1193–1197. <http://dx.doi.org/10.1086/515683>.
30. Bukh J, Meuleman P, Tellier R, Engle RE, Feinstone SM, Eder G, Satterfield WC, Govindarajan S, Krawczynski K, Miller RH, Leroux-Roels G, Purcell RH. 2010. Challenge pools of hepatitis C virus genotypes 1–6 prototype strains: replication fitness and pathogenicity in chimpanzees and human liver-chimeric mouse models. *J Infect Dis* 201:1381–1389. <http://dx.doi.org/10.1086/651579>.
31. Feinstone SM, Alter HJ, Dienes HP, Shimizu Y, Popper H, Blackmore D, Sly D, London WT, Purcell RH. 1981. Non-A, non-B hepatitis in chimpanzees and marmosets. *J Infect Dis* 144:588–598. <http://dx.doi.org/10.1093/infdis/144.6.588>.
32. Engle RE, Russell RS, Purcell RH, Bukh J. 2008. Development of a TaqMan assay for the six major genotypes of hepatitis C virus: comparison with commercial assays. *J Med Virol* 80:72–79. <http://dx.doi.org/10.1002/jmv.21043>.
33. Yanagi M, Bukh J, Emerson SU, Purcell RH. 1996. Contamination of commercially available fetal bovine sera with bovine viral diarrhea virus genomes: implications for the study of hepatitis C virus in cell cultures. *J Infect Dis* 174:1324–1327. <http://dx.doi.org/10.1093/infdis/174.6.1324>.
34. Corbet S, Bukh J, Heinsen A, Fomsgaard A. 2003. Hepatitis C virus subtyping by a core-envelope 1-based reverse transcriptase PCR assay with sequencing and its use in determining subtype distribution among Danish patients. *J Clin Microbiol* 41:1091–1100. <http://dx.doi.org/10.1128/JCM.41.3.1091-1100.2003>.
35. Rollier C, Depla E, Drexhage JA, Verschoor EJ, Verstrepen BE, Fatmi A, Brinster C, Fournillier A, Whelan JA, Whelan M, Jacobs D, Maertens G, Inchauste G, Heeney JL. 2004. Control of heterologous hepatitis C virus infection in chimpanzees is associated with the quality of vaccine-induced peripheral T-helper immune response. *J Virol* 78:187–196. <http://dx.doi.org/10.1128/JVI.78.1.187-196.2004>.
36. Yanagi M, Purcell RH, Emerson SU, Bukh J. 1999. Hepatitis C virus: an infectious molecular clone of a second major genotype (2a) and lack of viability of intertypic 1a and 2a chimeras. *Virology* 262:250–263. <http://dx.doi.org/10.1006/viro.1999.9889>.
37. Sakai A, Takikawa S, Thimme R, Meunier JC, Spangenberg HC, Govindarajan S, Farci P, Emerson SU, Chisari FV, Purcell RH, Bukh J. 2007. In vivo study of the HC-TN strain of hepatitis C virus recovered from a patient with fulminant hepatitis: RNA transcripts of a molecular clone (pHC-TN) are infectious in chimpanzees but not in Huh7.5 cells. *J Virol* 81:7208–7219. <http://dx.doi.org/10.1128/JVI.01774-06>.
38. Farci P, Shimoda A, Wong D, Cabezon T, De Gioannis D, Strazzer A, Shimizu Y, Shapiro M, Alter HJ, Purcell RH. 1996. Prevention of hepatitis C virus infection in chimpanzees by hyperimmune serum against the hypervariable region 1 of the envelope 2 protein. *Proc Natl Acad Sci U S A* 93:15394–15399. <http://dx.doi.org/10.1073/pnas.93.26.15394>.
39. Yanagi M, Purcell RH, Emerson SU, Bukh J. 1997. Transcripts from a single full-length cDNA clone of hepatitis C virus are infectious when directly transfected into the liver of a chimpanzee. *Proc Natl Acad Sci U S A* 94:8738–8743. <http://dx.doi.org/10.1073/pnas.94.16.8738>.
40. Jensen TB, Gottwein JM, Scheel TK, Hoegh AM, Eugen-Olsen J, Bukh J. 2008. Highly efficient JFH1-based cell-culture system for hepatitis C virus genotype 5a: failure of homologous neutralizing-antibody treatment to control infection. *J Infect Dis* 198:1756–1765. <http://dx.doi.org/10.1086/593021>.
41. Krawczynski K, Alter MJ, Tankersley DL, Beach M, Robertson BH, Lambert S, Kuo G, Spelbring JE, Meeks E, Sinha S, Carson DA. 1996. Effect of immune globulin on the prevention of experimental hepatitis C virus infection. *J Infect Dis* 173:822–828. <http://dx.doi.org/10.1093/infdis/173.4.822>.
42. Prentoe J, Jensen TB, Meuleman P, Serre SB, Scheel TK, Leroux-Roels G, Gottwein JM, Bukh J. 2011. Hypervariable region 1 differentially impacts viability of hepatitis C virus strains of genotypes 1 to 6 and impairs virus neutralization. *J Virol* 85:2224–2234. <http://dx.doi.org/10.1128/JVI.01594-10>.
43. Vanwolleghem T, Bukh J, Meuleman P, Desombere I, Meunier JC, Alter H, Purcell RH, Leroux-Roels G. 2008. Polyclonal immunoglobulins from a chronic hepatitis C virus patient protect human liver-chimeric mice from infection with a homologous hepatitis C virus strain. *Hepatology* 47:1846–1855. <http://dx.doi.org/10.1002/hep.22244>.
44. Morin TJ, Broering TJ, Leav BA, Blair BM, Rowley KJ, Boucher EN, Wang Y, Cheslock PS, Knauber M, Olsen DB, Ludmerer SW, Szabo G, Finberg RW, Purcell RH, Lanford RE, Ambrosino DM, Molrine DC, Babcock GJ. 2012. Human monoclonal antibody HCV1 effectively prevents and treats HCV infection in chimpanzees. *PLoS Pathog* 8:e1002895. <http://dx.doi.org/10.1371/journal.ppat.1002895>.
45. Pestka JM, Zeisel MB, Blaser E, Schurmann P, Bartosch B, Cosset FL, Patel AH, Meisel H, Baumert J, Viazov S, Rispeter K, Blum HE, Roggendorf M, Baumert TF. 2007. Rapid induction of virus-neutralizing antibodies and viral clearance in a single-source outbreak of hepatitis C. *Proc Natl Acad Sci U S A* 104:6025–6030. <http://dx.doi.org/10.1073/pnas.0607026104>.
46. Osburn WO, Snider AE, Wells BL, Latanich R, Bailey JR, Thomas DL, Cox AL, Ray SC. 2014. Clearance of hepatitis C infection is associated with the early appearance of broad neutralizing antibody responses. *Hepatology* 59:2140–2151. <http://dx.doi.org/10.1002/hep.27013>.
47. Falkowska E, Kajumo F, Garcia E, Reinus J, Dragic T. 2007. Hepatitis C virus envelope glycoprotein E2 glycans modulate entry, CD81 binding, and neutralization. *J Virol* 81:8072–8079. <http://dx.doi.org/10.1128/JVI.00459-07>.
48. Helle F, Goffard A, Morel V, Duverlie G, McKeating J, Keck ZY, Fong S, Penin F, Dubuisson J, Voisset C. 2007. The neutralizing activity of anti-hepatitis C virus antibodies is modulated by specific glycans on the E2 envelope protein. *J Virol* 81:8101–8111. <http://dx.doi.org/10.1128/JVI.00127-07>.
49. Pantua H, Diao J, Ultsch M, Hazen M, Mathieu M, McCutcheon K, Takeda K, Date S, Cheung TK, Phung Q, Hass P, Arnott D, Hongo JA, Matthews DJ, Brown A, Patel AH, Kelley RF, Eigenbrot C, Kapadia SB. 2013. Glycan shifting on hepatitis C virus (HCV) E2 glycoprotein is a mechanism for escape from broadly neutralizing antibodies. *J Mol Biol* 425:1899–1914. <http://dx.doi.org/10.1016/j.jmb.2013.02.025>.
50. Pedersen J, Carlsen TH, Prentoe J, Ramirez S, Jensen TB, Fornis X, Alter H, Fong SK, Law M, Gottwein J, Weis N, Bukh J. 2013. Neutralization resistance of hepatitis C virus can be overcome by recombinant human monoclonal antibodies. *Hepatology* 58:1587–1597. <http://dx.doi.org/10.1002/hep.26524>.
51. Farci P, Shimoda A, Coiana A, Diaz G, Peddis G, Melpolder JC, Strazzer A, Chien DY, Munoz SJ, Balestrieri A, Purcell RH, Alter HJ. 2000. The outcome of acute hepatitis C predicted by the evolution of the viral quasispecies. *Science* 288:339–344. <http://dx.doi.org/10.1126/science.288.5464.339>.
52. Giang E, Dornier M, Prentoe JC, Dreux M, Evans MJ, Bukh J, Rice CM, Ploss A, Burton DR, Law M. 2012. Human broadly neutralizing antibodies to the envelope glycoprotein complex of hepatitis C virus. *Proc Natl Acad Sci U S A* 109:6205–6210. <http://dx.doi.org/10.1073/pnas.1114927109>.
53. Keck Z, Wang W, Wang Y, Lau P, Carlsen TH, Prentoe J, Xia J, Patel AH, Bukh J, Fong SK. 2013. Cooperativity in virus neutralization by human monoclonal antibodies to two adjacent regions located at the amino terminus of hepatitis C virus E2 glycoprotein. *J Virol* 87:37–51. <http://dx.doi.org/10.1128/JVI.01941-12>.
54. Keck ZY, Xia J, Wang Y, Wang W, Krey T, Prentoe J, Carlsen T, Li AY, Patel AH, Lemon SM, Bukh J, Rey FA, Fong SK. 2012. Human monoclonal antibodies to a novel cluster of conformational epitopes on HCV E2 with resistance to neutralization escape in a genotype 2a isolate. *PLoS Pathog* 8:e1002653. <http://dx.doi.org/10.1371/journal.ppat.1002653>.